

BLADDER OUT OF CONTROL? WE CAN HELP



Overactive Bladder (OAB) is a treatable condition. It's not a normal part of aging. And you shouldn't have to deal with it on your own.

DOES THIS SOUND FAMILIAR?

- Going more than 8 times a day
- Avoiding social events
- Using pads to control leaks

OAB IS EXTREMELY COMMON

1 in 6 adults^{1,2} has OAB



VS.

21 million adults in the U.S. with vision problems³

29 million adults in the U.S. with diabetes⁴


IGNORING OAB POSES SERIOUS RISKS

People with undiagnosed OAB reported⁵:

54% Loss of confidence

49% Loss of self-esteem

45% Loss of intimacy



4.5 out of 10 adults **DON'T SEEK HELP** for overactive bladder⁵

IF OAB DISRUPTS YOUR LIFE, DON'T CHOOSE THE SILENT TREATMENT

Just say: "I think I might have OAB" and let your doctor take it from there.

FIND YOUR WAY TO REAL RELIEF

There are many ways to manage OAB. Remember, if conservative treatments don't deliver the results you need, you have more options.


DIAGNOSIS



"What is happening to me?"

Meet with a urinary incontinence specialist to see if you have OAB.

LIFESTYLE CHANGES



"What should I try first?"

Conservative treatments can help some people, but may not work well for others.

- Diet and exercise
- Pelvic floor strengthening (Kegels)
- Bladder training

ORAL MEDICATIONS



"Are there medications for OAB?"

Oral medications can help, but may cause side effects.

ADVANCED THERAPIES



"What if oral medications don't work?"

NEUROMODULATION

Therapies that use gentle nerve stimulation thought to normalize bladder-brain communication.

Medtronic Bladder Control Therapy delivered by the InterStim™ systems

- Try it during an evaluation
- Proven long-term relief⁶

Implanting an InterStim™ system has risks similar to any surgical procedure, including swelling, bruising, bleeding, and infection. Talk with your doctor about ways to minimize these risks.

Medtronic Bladder Control Therapy delivered by the NURO™ system

- In-office treatment

The most common side effects associated with the NURO™ system are temporary and include mild pain or skin inflammation at or near the stimulation site.

1. Stewart WF, et al. Prevalence and burden of overactive bladder in the United States. *World J Urol.* 2003 May;20(6):327-336.
2. United Nations, Department of Economic and Social Affairs, Population Division (2011). *World Population Prospects: The 2010 Revision, CD-ROM Edition.*
3. National Center for Health Statistics, National Health Interview Survey (NHIS). http://www.cdc.gov/visionhealth/basic_information/vision_loss_burden.htm. Accessed October 5, 2016.
4. National diabetes statistics, 2011. National Diabetes Information Clearinghouse website. <http://www.diabetes.niddk.nih.gov/dm/pubs/statistics/#fast>. Updated September 09, 2013. Accessed September 25, 2013.
5. Leede Research. "Views on OAB: A Study for the National Association of Continence." December 16, 2015.
6. Medtronic-sponsored research. InterStim Therapy Clinical Summary 2018.
7. Visco AG, Brubaker L, Richter HE, et al. Anticholinergic therapy vs. onabotulinumtoxinA for urgency urinary incontinence. *N Engl J Med.* 2012 Nov 8;367(19):1803-13.

INJECTED MEDICATIONS

Injections into the bladder repeated every 3-6 months.

- Potentially requires self-catheterization⁷
- Increases the risk of bladder infection due to self-catheterization⁷

TALK TO YOUR DOCTOR

or visit medtronic.com/bladder
to find a specialist today.



NATIONAL ASSOCIATION FOR CONTINENCE

Medtronic NURO™ Percutaneous Tibial Neuromodulation (PTNM) treats overactive bladder and associated symptoms of urinary urgency, urinary frequency, and urge incontinence. PTNM is not intended for patients with pacemakers or implantable defibrillators, patients prone to excessive bleeding, patients with nerve damage that could impact either percutaneous tibial nerve or pelvic floor function, or on patients who are pregnant or planning pregnancy. Do not use if the skin in the area of use is compromised. Exercise caution for patients with heart problems. Adverse events are typically temporary, and include mild pain, minor inflammation and bleeding near treatment site.

This therapy is not for everyone. Please consult your physician to decide whether PTNM is right for you. A prescription is required. For further information, please call Medtronic at 1-800-328-0810.

USA Rx Only Rev 0915

Medtronic Bladder Control Therapy delivered by the InterStim™ system treats urinary retention (inability to completely empty the bladder) and the symptoms of overactive bladder, including urinary urge incontinence (leakage) and significant symptoms of urgency-frequency. It should be used after you have tried other treatments such as medications and behavioral therapy and they have not worked, or you could not tolerate them.

You must demonstrate an appropriate response to the evaluation to be a candidate. You cannot have diathermy (deep heat treatment from electromagnetic energy) if you have an InterStim™ device.

This therapy is not intended for patients with a urinary blockage. Safety and effectiveness have not been established for pregnancy and delivery; patients under the age of 16; or for patients with neurological disease origins.

In addition to risks related to surgery, complications can include pain at the implant sites, new pain, infection, lead (thin wire) movement/migration, device problems, interactions with certain other devices or diagnostic equipment such as MRI, undesirable changes in urinary or bowel function, and uncomfortable stimulation (sometimes described as a jolting or shocking feeling).

This therapy is not for everyone. This treatment is prescribed by your doctor. Please talk to your doctor to decide whether this therapy is right for you. Your doctor should discuss all potential benefits and risks with you. Although many patients may benefit from the use of this treatment, results may vary. For further information, please call Medtronic at 1-800-328-0810 and/or consult Medtronic's website at www.medtronic.com.

USA Rx Only. Rev 0517